

Biological and molecular characteristics of Cherry leaf roll virus (CLRV) isolates from different host plants

J Gentkow, K Rebenstorf, S von Bargaen, C Büttner

Humboldt-Universität zu Berlin, Institute for Horticultural Sciences, Berlin, Germany

Email: susanne.von.bargaen@agrar.hu-berlin.de

INTRODUCTION

Cherry leaf roll nepovirus (CLRV) is a widespread pathogen of woody plants in Germany and throughout the European Community (Bandte & Büttner, 2001). It is also spread in several herbaceous plants, such as rhubarb (*Rheum rhabarbarum*), but the origin of this virus is unclear (Jones, 1985). CLRV often induces symptoms in, for example, ash, birch, cherry, elderberry and walnut, including delayed leaf development, chlorotic leaf streaks or spots as well as dieback of branches or whole trees (Hamacher & Giersiepen, 1989; Hamacher & Quadt, 1991; Rebenstorf & Obermeier, 2003). In nature, CLRV is mainly transmitted through seed or pollen, leading to viral spread within one plant species; mechanical transmission, including grafting procedures, is also possible (Cooper *et al.*, 1984; Massalski & Cooper, 1984; Mircetich *et al.*, 1980). Genomic organisation of CLRV is still unknown, but morphology and particle composition is typical for nepoviruses, which are classified within the family of *Comoviridae* (Jones, 1985). Supporting the observation that CLRV isolates, obtained from identical host species, are serologically related, a phylogenetic analysis of a 280 base pair fragment of the 3' non-coding-region of viral RNA demonstrated that CLRV isolates were also clustered on a genomic level that accorded with the original host plant.

METHODS AND RESULTS

One-year-old seedlings of nine different plant species were mechanically inoculated and cultured in the field (Berlin, Germany), to evaluate the role of transmission via wounding as a source of horizontal virus spread between different plant species under natural environmental conditions. Ten selected CLRV isolates, originating from different host plants, were used for inoculation. Six months after inoculation no characteristic symptoms of CLRV infection were observed. However, buds from inoculated trees were sampled and analysed for CLRV infection by a sensitive IC-RT-PCR assay with specific primers, according to Werner *et al.* (1997). In none of the samples could an infection be confirmed. Furthermore, virus particles were purified after their propagation on the herbaceous host plant *Chenopodium quinoa*, in order to compare viral components (structure-proteins and nucleic acids) of CLRV isolates included in the field experiment. On *C. quinoa* all ten CLRV isolates induced chlorotic spots on inoculated leaves 3–5 days after inoculation. A systemic infection, exhibiting leaf distortion and necrosis of expanding leaves, was established 9 days after inoculation in all cases, with the exception of CLRV-walnut strain from Hungary; this particular isolate induced necrotic local lesions on inoculated *C. quinoa* leaves without systemic symptoms. Electron microscopic control of purified virus preparations revealed isometric virus particles (28 nm in diameter) typical for nepoviruses in all investigated isolates. Protein analysis of purified viruses by SDS-PAGE displayed, from eight CLRV

isolates, a structure-protein of expected size (MW is approximately 53 kDa). In two isolate purifications different proteins of significantly smaller size were present, suggesting an infection with another virus. In RNA-gels, seven virus isolates showed typical RNA patterns, with a larger genomic RNA1 (molecular weight around 8,2 kb) and a slightly shorter RNA2 (c. 6.9–7.3 kb). While the virus isolate from rhubarb (Germany) exhibited a 53 kDa coat protein typical for CLRV, only one RNA was detectable in RNA gels under native conditions. The CLRV isolate derived from European ash (Germany) as well as the CLRV-walnut strain from Hungary, which were different in protein analysis, also exhibited atypical RNA patterns. Correspondingly CLRV detection by RT-PCR, using particle purifications of these two isolates as templates, also failed.

To date, it remains unclear whether or not, in certain woody host plants, CLRV isolates appear as mixed infections with other viruses of similar morphology.

Further investigations are necessary to identify the viruses responsible for unusual RNA and protein patterns. Additionally, detection of CLRV in the initiated field experiments has to be continued to clarify the potential of different virus isolates to infect various woody host plants through wounds and the relevance of CLRV distribution between species by other factors than through seed or pollen.

REFERENCES

- Bandte M; Büttner C (2001). Occurrence, transmission and diagnosis of cherry leafroll nepovirus – a literature review. *Pflanzenschutzberichte* **59**, 1-19.
- Cooper JI; Massalski PR; Edwards ML (1984). Cherry leaf roll virus in the female gametophyte and seed of birch and its relevance to vertical transmission. *Annals of Applied Biology* **105**, 55-64.
- Hamacher J; Giersiepen R (1989). Histologisch-cytologische Veränderungen in Kirschenblattrollvirus-infizierten und durch Luftschadstoffe gestreßten Betula-Arten. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes* **41**, 124-130.
- Hamacher J; Quadt A (1991). Light- and Electron Microscopic Studies of Cherry Leaf Roll Virus (CLRV) on European Ash (*Fraxinus excelsior* L.). *Journal of Phytopathology* **131**, 215-226.
- Jones AT (1985) *Cherry leaf roll virus*. CM/AAB Description of plant viruses, No. 306.
- Massalski PR; Cooper JI (1984). The location of virus-like particles in the male gametophyte of birch, walnut, cherry naturally infected with CLRV and its relevance to vertical transmission of the virus. *Plant Pathology* **33**, 255-262.
- Mircetich SM; Sanborn RR; Ramos, DE (1980). Natural spread, graft-transmission, and possible ethiology of walnut blackline disease. *Phytopathology* **72**, 988.
- Rebenstorf K; Obermeier C (2003). Gefahr für Gehölze: Das Kirschenblattrollvirus. *Deutsche Baumschule* **7**, 54-55.
- Werner R; Mühlbach, HP; Büttner C (1997). Detection of cherry leaf roll nepovirus (CLRV) in birch, beech and petunia by immunocapture RT-PCR using a conserved primerpair. *European Journal for Pathology* **27**, 309-318.